# HART BEAT

19<sup>th</sup> July 2017 ISSUE 42 Hart Field Site Group Inc

PO Box 939 CLARE SA 5453

0427 423 154

admin@hartfieldsite.org.au







### IN THIS ISSUE

### **Definitions**

Site information

### Hart Beat site reports

- Hart
- Spalding
- Condowie
- Kybunga
- Farrell Flat
- Pinery
- Eudunda
- Tarlee

**NOTE** - there is large variation in yield predictions among sites this season. This can be attributed to the uncertainly of seasonal conditions and rainfall for the remainder of the 2017 growing season.

### HART BEAT

# Yield Prophet® simulations for 8 sites across the mid-north of SA

The Yield Prophet<sup>®</sup> simulations featured are not a crystal ball, but provide a realistic prediction of the available soil water and nitrogen status of your crop



## **HART EVENTS**

HART FIELD DAY – 19<sup>th</sup> September 2017 Spring Twilight Walk – 17<sup>th</sup> October 2017 Getting The Crop In – March 2018

# **Definitions**

### Hart Beat definitions

Each site has been characterised for plant available water capacity (PAWC) and bulk density to determine how much of the measured water and nitrogen is available to the crop during the season.

**Drained upper limit (DUL)** – is the amount of water that a saturated soil holds after it has drained.

**Crop lower limit (CLL)** – is the amount of water remaining in the soil after crop senescence.

Plant available water capacity (PAWC) – is the difference between the drained upper limit of the soil and the lower extraction limit of a crop over the depth of rooting. It is the maximum water available to a crop from a particular soil type.

**Plant available water (PAW)** – is the amount of water contained in the soil at a given time minus the crop lower limit.

**Bulk density (BD)** – is a measure of the weight of dry soil per unit volume of soil.

**Growing season rainfall (GSR)** – is rainfall for the period between and including April to October.

**Decile** – is a measure of seasonal rainfall on a scale of 1 to 9. In a decile 7 year, 70% of previous years were dryer, in a decile 3 year 30% of previous years were dryer.

Yield probability curves - display two different nitrogen scenarios for each site. The green line displays the actual grain yield with the current soil available nitrogen. The blue line represents the grain yield potential with unlimited nitrogen (yield potential). A small difference between these two lines indicates the current soil N level is adequate for the crop to reach its yield potential. Conversely, a large difference between these two lines indicates

additional N fertiliser is required for the crop to reach its yield potential.

The **French & Schultz** formula estimates the rainfall limited grain yield based on the growing season rainfall (GSR). It assumes evaporation of 110mm, includes stored water at sowing (30% of Jan to Mar rainfall) and a maximum grain yield potential of 20 kg/mm/ha.

Yield Potential = GSR (Apr-Oct) – Evaporation (110mm) \* 20 kg/mm/ha.

**Yield Prophet**® has been very accurate throughout Australia, over the past 5 seasons. At the Hart field site the *Yield Prophet*® prediction on the 15<sup>th</sup> September, using an average finish, has been only 16% above the final grain yield, averaged over the past 4 years, making wheat growth models such as APSIM highly valuable.

Yield Prophet® is an internet based service which uses the APSIM wheat prediction model.

The model relies on accurate soil, crop, historical climate data and up to date local weather information to predict plant growth rates and final hay or grain yields. These are critical measurements specific to the site being analysed and may not fit closely to individual situations. Instead the predictions will give a realistic guide to seasonal prospects based on a site with similar rainfall and / or soil type.

Using climate data for the current season, *Yield Prophet*<sup>®</sup> simulates the soil water and nitrogen processes in the paddock, and crop growth. *Yield Prophet*<sup>®</sup> calculates the amount of water and nitrogen available to the crop and the water and nitrogen demand of the crop.

**Disclaimer:** Yield Prophet® information is used entirely at your own risk. You will accept all risks and responsibility for losses, damages, costs and other consequences of using Yield Prophet® information and reports. To the maximum extent permitted by law, APSRU and BCG excludes all responsibility and liability to any person arising directly or indirectly from using the information generated by Yield Prophet®.

**Important Notice:** Yield Prophet® does not generate recommendations or advice, it is only a guide and must be combined with local paddock and district knowledge. APSIM does not take into account weed competition, pest/disease pressure, pesticide / herbicide damage, farmer error, or extreme events (such as extreme weather, flood and fire). For more information about APSIM or *Yield Prophet*® please visit or www.yieldprophet.com.au.

# Site information

### Rainfall and soil water characteristics for all sites

Site	Average annual rainfall (mm)	Soil type	Plant Available Water Capacity (mm)	Soil Profile Depth (cm)	Plant Available Water at 10 May* (mm)	Pre-sowing soil nitrogen (kg/ha)
Hart	400	Sandy clay loam	206	150	92	90
Spalding	430	Red brown earth	143	150	57	90
Condowie	350	Sandy loam	115	150	40	90
Kybunga	428	Clay loam	262	120	89	88
Farrell Flat	474	Light clay loam	172	120	63	87
Pinery	374	Silty clay loam	79	150	43	90
Eudunda	445	Gravelly loam	96	100	0	90
Tarlee	474	Sandy loam	113	150	66	87

<sup>\*</sup>Note there were errors in the input of PAW values in the June 2 Hart Beat.

### 2017 site locations



## www.hartfieldsite.org.au

### SANDY CLAY LOAM

**Crop growth** 

Variety: Mace wheat Sowing date: 1st May Nitrogen fertiliser: 40 kg N/ha at seeding + 20 kg N/ha July 17

The season so far

Annual rain to date: 176 mm

GSR to date: 105 mm (23 mm since last report) GSR decile: 2
Current predicted PAW: 61 mm (30 % full) PAWC: 206 mm

#### **Grain yield predictions (Yield Prophet)**

**Yield prophet estimate:** (Date of report 19/07/2017)


These estimates are based on a 50% probability

Yield t/ha	Sown 1st May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	4.3	- 0.5	4.3	- 0.8

#### French & Schultz grain yield estimate:


100% WUE:	3.1 <b>t/ha</b>
80% WUE:	2.5 t/ha

This model assumes that there is 21 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (141 mm) for the rest of the season.



The graphs above and below show the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

#### Hay Yield Outcome



### **SPALDING**

### **RED BROWN EARTH**

Crop growth

Variety: Mace wheat Sowing date: Nitrogen fertiliser: 1st May 40 kg N/ha at seeding

The season so far

Annual rain to date: 184mm

GSR to date: 115 mm (27 mm since last report)

32 mm (23% full) **Current predicted PAW:** 

**Grain yield predictions (Yield Prophet)** 

Yield prophet estimate: (Date of report 19/07/2017)

These estimates are based on a 50% probability

Yield t/ha	Sown 1 <sup>st</sup> May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	3.6	- 0.7	3.2	- 0.4


#### French & Schultz grain yield estimate:

100% WUE:	3.1 t/ha
80% WUE:	2.5 t/ha

This model assumes that there is 21 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (130 mm) for the rest of the season.

**GSR** decile: 2 PAWC: 143 mm

#### Grain yield outcome graph



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

## **CONDOWIE**

### **SANDY LOAM**

Crop growth

Variety: Mace wheat Nitrogen fertiliser: 40 kg N/ha at seeding Sowing date: 1st May

The season so far

Annual rain to date: 181 mm

GSR to date: 101 mm (25 mm since last report)

**Current predicted PAW:** 52 mm (45 % full)

#### **Grain yield predictions (Yield Prophet)**

Yield prophet estimate: (Date of report 11/07/2017)

These estimates are based on a 50% probability

Yield t/ha	Sown 1st May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	2.6	- 0.9	2.5	- 0.2

#### French & Schultz grain yield estimate:

100% WUE:	2.6 t/ha
80% WUE:	2.1 t/ha

This model assumes that there is 24 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (114 mm) for the rest of the season.


### Grain yield outcome graph

3

115 mm

**GSR** decile:

PAWC:



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

### **KYBUNGA**

### **CLAY LOAM**

Crop growth

Variety: Mace wheat Sowing date: 1st May Nitrogen fertiliser: 40 kg N/ha at seeding + 20 kg N/ha July 17

The season so far

Annual rain to date: 210 mm

GSR to date: 102 mm (36 mm since last report)

**Current predicted PAW:** 86 mm (33 % full)

**Grain yield predictions (Yield Prophet)** 

Yield prophet estimate: (Date of report 11/07/2017)

These estimates are based on a 50% probability

Yield t/ha	Sown 1 <sup>st</sup> May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	4.8	- 0.2	4.6	- 0.8

#### French & Schultz grain yield estimate:


100% WUE:	3.7 t/ha
80% WUE:	2.9 t/ha

This model assumes that there is 35 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (159 mm) for the rest of the season.

**GSR** decile:

PAWC: 262 mm

#### Grain yield outcome graph



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

### **FARRELL FLAT**

### LIGHT CLAY LOAM

**Crop growth** 

Variety: Mace wheat Sowing date: Nitrogen fertiliser: 40 kg N/ha at seeding + 20 kg N/ha July 17 1st May

The season so far

Annual rain to date: 194 mm

**GSR** to date: 111 mm (49 mm since last report) **GSR** decile: PAWC: 172 mm

**Current predicted PAW:** 38 mm (22% full)

#### **Grain yield predictions (Yield Prophet)**

(Date of report 19/07/2017) Yield prophet estimate:

These estimates are based on a 50% probability


Yield t/ha	Sown 1st May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	4.7	-0.3	4.5	- 0.6

#### French & Schultz grain yield estimate:

100% WUE:	3.7 t/ha
80% WUE:	3.0 t/ha

This model assumes that there is 25 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (160 mm) for the rest of the season.

#### Grain yield outcome graph



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

### **SILTY CLAY LOAM**

Crop growth

Variety: Mace wheat Sowing date: Nitrogen fertiliser: 40 kg N/ha at seeding 1st May

The season so far

Annual rain to date: 203 mm

GSR to date: 83 mm (29 mm since last report)

**Current predicted PAW:** 25 mm (32 % full)

**Grain yield predictions (Yield Prophet)** 

Yield prophet estimate: (Date of report 19/07/2017)

These estimates are based on a 50% probability


Yield t/ha	Sown 1 <sup>st</sup> May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	3.8	- 0.4	3.5	- 0.6

#### French & Schultz grain yield estimate:

100% WUE:	2.8 t/ha
80% WUE:	2.2 t/ha

This model assumes that there is 36 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (129 mm) for the rest of the season. **GSR** decile: 1 PAWC: 79 mm

#### Grain yield outcome graph



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

### **EUDUNDA**

### **GRAVELLY LOAM**

Crop growth

Variety: Mace wheat Sowing date: Nitrogen fertiliser: 40 kg N/ha at seeding 1st May

The season so far

Annual rain to date: 209 mm

**GSR** to date: 117 mm (17 mm since last report)

**Current predicted PAW:** 14 mm (15 % full)

#### **Grain yield predictions (Yield Prophet)**

Yield prophet estimate: (Date of report 19/07/2017)

These estimates are based on a 50% probability

	Yield t/ha	Sown 1st May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
ĺ	Grain	2.8	- 0.3	3.1	- 0.1

#### French & Schultz grain yield estimate:

100% WUE:	3.7 t/ha
80% WUE:	2.9 t/ha
OU% WUE.	Z.9 VIId


This model assumes that there is 28 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (149 mm) for the rest of the season.

### Grain yield outcome graph

96 mm

**GSR** decile:

PAWC:



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

### **SANDY LOAM**

**Crop growth** 

Variety: Mace wheat Sowing date: 1st May Nitrogen fertiliser: 40 kg N/ha at seeding + 20 kg N/ha July 17

The season so far

**Annual rain to date:** 181 mm

**GSR to date:** 87 mm (34 mm since last report)

Current predicted PAW: 54 mm (48 % full)

**Grain yield predictions (Yield Prophet)** 

**Yield prophet estimate:** (Date of report 19/07/2017)

These estimates are based on a 50% probability

Yield t/ha	Sown 1 <sup>st</sup> May (see graph)	Change Since Last Report	Sown 20 <sup>th</sup> May	Change Since Last Report
Grain	4.5	+ 0.3	4.9	+ 0.5


#### French & Schultz grain yield estimate:

100% WUE:	3.5 t/ha
80% WUE:	2.8 t/ha

This model assumes that there is 28 mm stored moisture, 110 mm of evaporation and Decile 5 rainfall (169 mm) for the rest of the season.

GSR decile: 1 PAWC: 113 mm

#### Grain yield outcome graph



This graph shows the chance of reaching the corresponding yield given weather, soil conditions and agronomic inputs to date, and historical climate data (100yrs) to simulate remainder of the season.

## Around the site



Photos taken at Hart on July 19, 2017



